2.5 cm

2.5 cm

2.5 cm

2.5 cm

STUDY OF THE FLUORINE AND BORON-10 CONTAINING COMPOUNDS TOWARD MRI AND

12 pt. bold ALL CAPS

Leave a blank line (18 pt. fixed).

Yoshihide Hattori[1], Yoshihiro Yamaguchi[1], Hitoshi Yamamoto[2], Tomohiro Asano[3], Mitsunori Kiriha[4], Hiteaki Wakamiya[1]

Underline for a speaker

Leave a blank line (18 pt. fixed).

[1]Faculty of Science and Technology, Kinki University, Higashi-osaka, Osaka 577-8502, Japan, [2]Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan, [3]College of Agriculture, Univer Osaka 599-8231, Japan, [4]Department of Neurosurgery, Aino Junior College Hospital, Ibaraki, Osaka 567-0018, Japan

12 pt. (18 pt. single-spaced)

Leave a blank line (18 pt. fixed).

Dipeptides containing 3-(4-fluorophenyl)alanine [Phe(F)] (**1**) seem to be transferred into some kinds of tumor cells through the oligopeptide transporter. Furthermore, in a previous stud ining 3-(2,3,4,5,6-pentafluorophenyl)alanine [Phe(F_5)] (**2**) was certified to be detectable by ^{19}F NMR up to μM order concentration. These facts suggest that magnetic resonance imaging (M t of the Phe(F_5)-containing peptides internali promising means for diagnosis of cancer.

No subtitles

12 pt. typeface with 1.5-line spacing (18 pt. fixed).

From the standpoint of the treatment of brain cancer or melanoma, the boron neutron capture therapy (BNCT) based on the interaction of ^{10}B isotope and neutron has been highly noted in recent years [1]. In l and BNCT, we designed and synthesized the n and boron-10 atoms such as 3-(4-borono B] (**3**) and 3-(4-borono-2,6-difluorophenyl)alaninol {[Bpa(F_2)-^{10}B]-ol} (**4**). In the present paper we focus on ^{19}F NMR measurement and tumor cell killing effect of various compounds containing both fluorine and boron-10 atoms.

Leave a blank line (18 pt. fixed).



Phe(F) (**1**) Phe(F_5) (**2**) Bpa(F_2)-^{10}B (**3**) Bpa(F_2)-ol-^{10}B (**4**)

Leave a blank line (18 pt. fixed).

[1] Soloway, H. A., Tjarks, W., Barnum, A. B., Rong, F., Barth, F. R., Codogni, M. I., Wilson. J. G. (1998) *Chem. Rev.*, **98**, 151 1562.

12 pt. (18 pt. single-spaced)

3.0 cm